

Michigan Department of Health and Human Services Preventive Services Coverage Guidelines

The following is a list of recommended preventive services and corresponding CPT, HCPCS, and diagnosis codes to assist Medicaid health plans and providers in identifying Medicaid covered preventive services. Providers are encouraged to identify the service as preventive by including HCPCS modifier 33, preventive service. In addition to modifier 33, a diagnosis may be required to support that the service was preventive.

This information is provided as a reference only and not intended to replace the clinical judgement of the providers. Other coverage determinations may apply.

Additional information about preventive care guidelines is available at the following:

- United States Preventive Services Task Force grade A and B services
<http://www.uspreventiveservicestaskforce.org/Page/Name/recommendations>
- Advisory Committee on Immunization Practices recommended vaccines
<http://www.cdc.gov/vaccines/hcp/acip-recs/index.html>
- Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine recommended preventive services for women
<http://www.nationalacademies.org/hmd/Reports/2011/Clinical-Preventive-Services-for-Women-Closing-the-Gaps.aspx>
- Early and Periodic Screening, Diagnosis and Treatment services as defined in the current periodicity schedule by the American Academy of Pediatrics
http://brightfutures.aap.org/clinical_practice.html

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Abdominal Aortic Aneurysm: Screening: men aged 65 to 75 years who have ever smoked	The USPSTF recommends 1-time screening for abdominal aortic aneurysm (AAA) with ultrasonography in men aged 65 to 75 years who have ever smoked.	76706, 76770*, 76775*	Z136, Z87891
Abnormal Blood Glucose and Type 2 Diabetes Mellitus: Screening: adults aged 40 to 70 years who are overweight or obese	The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity.	82947*, 82948*, 82950*, 82951*, 82952*, 83036*	Z131, Z833

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Aspirin Use to Prevent Cardiovascular Disease and Colorectal Cancer: Preventive Medication: adults aged 50 to 59 years with a 10% or greater 10-year cvd risk	The USPSTF recommends initiating low-dose aspirin use for the primary prevention of cardiovascular disease (CVD) and colorectal cancer (CRC) in adults aged 50 to 59 years who have a 10% or greater 10-year CVD risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.	Pharmacy Benefit	NA
Asymptomatic Bacteriuria in Adults: Screening: pregnant persons	The USPSTF recommends screening for asymptomatic bacteriuria using urine culture in pregnant persons.	87086*, 87088*	Reason For Screen
BRCA-Related Cancer: Risk Assessment, Genetic Counseling, and Genetic Testing: women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or an ancestry associated with brca1/2 gene mutation	The USPSTF recommends that primary care clinicians assess women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or who have an ancestry associated with breast cancer susceptibility 1 and 2 (BRCA1/2) gene mutations with an appropriate brief familial risk assessment tool. Women with a positive result on the risk assessment tool should receive genetic counseling and, if indicated after counseling, genetic testing.	Assessment and counseling covered during preventive medicine services or a focused E/M visit. Labs as indicated. 81162-81167, 81212-81217	Reason For Visit
Breast Cancer: Medication Use to Reduce Risk: women at increased risk for breast cancer aged 35 years or older	The USPSTF recommends that clinicians offer to prescribe risk-reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer and at low risk for adverse medication effects.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Breast Cancer: Screening: women aged 50 to 74 years	The USPSTF recommends biennial screening mammography for women aged 50 to 74 years.	77063, 77067	Reason For Screen
Breastfeeding: Primary Care Interventions: pregnant women, new mothers, and their children	The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	S9443	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Cervical Cancer: Screening: women aged 21 to 65 years	The USPSTF recommends screening for cervical cancer every 3 years with cervical cytology alone in women aged 21 to 29 years. For women aged 30 to 65 years, the USPSTF recommends screening every 3 years with cervical cytology alone, every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or every 5 years with hrHPV testing in combination with cytology (cotesting). See the Clinical Considerations section for the relative benefits and harms of alternative screening strategies for women 21 years or older.	87623-87625, 88141-88148, 88155, 88164-88167, 88174-88175, G0101, G0476, Q0091	Reason For Screen
Chlamydia and Gonorrhea: Screening: sexually active women	The USPSTF recommends screening for chlamydia in sexually active women age 24 years and younger and in older women who are at increased risk for infection.	87110, 87270, 87320, 87490-87492, 87810	Reason For Screen
Chlamydia and Gonorrhea: Screening: sexually active women	The USPSTF recommends screening for gonorrhea in sexually active women age 24 years and younger and in older women who are at increased risk for infection.	87590, 87591, 87592, 87850	Reason For Screen
Colorectal Cancer: Screening: adults aged 45 to 75 years	The USPSTF recommends screening for colorectal cancer in all adults aged 45 to 75 years. See the "Practice Considerations" section and Table 1 for details about screening strategies.	00812, 45330*, 45331*, 45333*, 45338*, 45346*, 45378*, 45380, 45384*, 45385*, 45388*, 81528, 82270, 82274*, G0104, G0105, G0121, G0328,	Z1211, Z1212, Z1213, Z800, Z8371, Z86.004, Z86010
Dental Caries in Children from Birth Through Age 5 Years: Screening: children from birth through age 5 years	The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption. The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is deficient in fluoride.	99188, D1206, D1208	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Depression in Adults: Screening: general adult population, including pregnant and postpartum women	The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	96127*	Reason For Visit
Depression in Children and Adolescents: Screening: adolescents aged 12 to 18 years	The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	96127	Reason For Visit
Falls Prevention in Community-Dwelling Older Adults: Interventions: adults 65 years or older	The USPSTF recommends exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Folic Acid for the Prevention of Neural Tube Defects: Preventive Medication: women who are planning or capable of pregnancy	The USPSTF recommends that all women who are planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.	Pharmacy Benefit	NA
Gestational Diabetes Mellitus, Screening: asymptomatic pregnant women, after 24 weeks of gestation	The USPSTF recommends screening for gestational diabetes mellitus (GDM) in asymptomatic pregnant women after 24 weeks of gestation.	82950*, 82951*, 82952*	Reason For Screen
Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults With Cardiovascular Risk Factors: Behavioral Counseling Interventions: adults with cardiovascular disease risk factors	The USPSTF recommends offering or referring adults with cardiovascular disease risk factors to behavioral counseling interventions to promote a healthy diet and physical activity.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Hearing loss screening: newborns	The USPSTF recommends screening for hearing loss in all newborn infants.	Inpatient Newborn Screen 92551-92553, 92558, 92568, 92650-92653, 92588, V5008	Reason For Visit
Hemoglobinopathies screening: newborns	The USPSTF recommends screening for sickle cell disease in newborns.	Inpatient Newborn Screen	Reason For Visit
Hepatitis B Virus Infection in Pregnant Women: Screening: pregnant women	The USPSTF recommends screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit	80055,80081, 86704-86706, 87340, 87341, G0499	Reason For Screen
Hepatitis C Virus Infection in Adolescents and Adults: Screening: adults aged 18 to 79 years	The USPSTF recommends screening for hepatitis C virus (HCV) infection in adults aged 18 to 79 years.	86803, G0472	Reason For Screen
Human Immunodeficiency Virus (HIV) Infection: Screening: adolescents and adults aged 15 to 65 years	The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults aged 15 to 65 years. Younger adolescents and older adults who are at increased risk of infection should also be screened. See the Clinical Considerations section for more information about assessment of risk, screening intervals, and rescreening in pregnancy.	86689*, 86701*, 86702*, 86703*, 87389*, 87390*, 87391*, 87534*, 87535*, 87536*, 87537*, 87538*, 87539*, 87806*, G0432, G0433, G0435	Z114
Human Immunodeficiency Virus (HIV) Infection: Screening: pregnant persons	The USPSTF recommends that clinicians screen for HIV infection in all pregnant persons, including those who present in labor or at delivery whose HIV status is unknown.	80081*, 86689*, 86701*, 86702*, 86703*, 87389*, 87390*, 87391*, 87534*, 87535*, 87536*, 87537*, 87538*, 87539*, 87806*, G0432, G0433, G0435	Reason For Screen
Hypothyroidism screening: newborns	The USPSTF recommends screening for congenital hypothyroidism in newborns.	Inpatient Newborn Screen	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Interventions for Tobacco Smoking Cessation in Adults, Including Pregnant Persons: nonpregnant adults	The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and US Food and Drug Administration (FDA)--approved pharmacotherapy for cessation to nonpregnant adults who use tobacco.	99406, 99407	Reason For Visit
Interventions for Tobacco Smoking Cessation in Adults, Including Pregnant Persons: pregnant persons	The USPSTF recommends that clinicians ask all pregnant persons about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant persons who use tobacco.	99406, 99407	Reason For Visit
Intimate Partner Violence, Elder Abuse, and Abuse of Vulnerable Adults: Screening: women of reproductive age	The USPSTF recommends that clinicians screen for intimate partner violence (IPV) in women of reproductive age and provide or refer women who screen positive to ongoing support services. See the Clinical Considerations section for more information on effective ongoing support services for IPV and for information on IPV in men.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Latent Tuberculosis Infection: Screening: asymptomatic adults at increased risk for infection	The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk.	86480*, 86481*, 86580*, 87116*	Z111, Z201
Low-Dose Aspirin Use for the Prevention of Morbidity and Mortality From Preeclampsia: Preventive Medication : pregnant women who are at high risk for preeclampsia	The USPSTF recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.	Pharmacy Benefit	NA
Lung Cancer: Screening: adults aged 50 to 80 years who have a 20 pack-year smoking history and currently smoke or have quit within the past 15 years	The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 50 to 80 years who have a 20 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	71250*, 71271*	F17200-F17299, Z122, Z720, Z87891,

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Obesity in Children and Adolescents: Screening: children and adolescents 6 years and older	The USPSTF recommends that clinicians screen for obesity in children and adolescents 6 years and older and offer or refer them to comprehensive, intensive behavioral interventions to promote improvements in weight status.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Ocular Prophylaxis for Gonococcal Ophthalmia Neonatorum: Preventive Medication: newborns	The USPSTF recommends prophylactic ocular topical medication for all newborns to prevent gonococcal ophthalmia neonatorum.	Newborn Inpatient	Reason For Visit
Osteoporosis to Prevent Fractures: Screening: postmenopausal women younger than 65 years at increased risk of osteoporosis	The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool. See the Clinical Considerations section for information on risk assessment.	76977*, 77078*, 77080*, 77081*	Z13820, Z8262
Osteoporosis to Prevent Fractures: Screening: women 65 years and older	The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 years and older.	76977*, 77078*, 77080*, 77081*	Z13820, Z8262
Perinatal Depression: Preventive Interventions: pregnant and postpartum persons	The USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal depression to counseling interventions.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Phenylketonuria screening: newborns	The USPSTF recommends screening for phenylketonuria in newborns.	Inpatient Newborn Screen	Reason For Visit
Preeclampsia: Screening: pregnant woman	The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Prevention and Cessation of Tobacco Use in Children and Adolescents: Primary Care Interventions: school-aged children and adolescents who have not started to use tobacco	The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents.	99406, 99407	Reason For Visit
Prevention of Human Immunodeficiency Virus (HIV) Infection: Preexposure Prophylaxis: persons at high risk of hiv acquisition	The USPSTF recommends that clinicians offer preexposure prophylaxis (PrEP) with effective antiretroviral therapy to persons who are at high risk of HIV acquisition. See the Clinical Considerations section for information about identification of persons at high risk and selection of effective antiretroviral therapy.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Rh(D) Incompatibility: Screening: pregnant women, during the first pregnancy-related care visit	The USPSTF strongly recommends Rh(D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.	80055*, 80081*, 86901*	Reason For Screen
Rh(D) Incompatibility: Screening: unsensitized rh(d)-negative pregnant women	The USPSTF recommends repeated Rh(D) antibody testing for all unsensitized Rh(D)-negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh(D)-negative.	80055*, 80081*, 86901*	Reason For Screen
Screening for Hepatitis B Virus Infection in Adolescents and Adults: adolescents and adults at increased risk for infection	The USPSTF recommends screening for hepatitis B virus (HBV) infection in adolescents and adults at increased risk for infection. See the Practice Considerations section for a description of adolescents and adults at increased risk for infection.	86704-86706, 87340, 87341, G0499	Reason For Screen
Screening for Hypertension in Adults: adults 18 years or older without known hypertension	The USPSTF recommends screening for hypertension in adults 18 years or older with office blood pressure measurement (OBPM). The USPSTF recommends obtaining blood pressure measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Sexually Transmitted Infections: Behavioral Counseling: sexually active adolescents and adults at increased risk	The USPSTF recommends behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs). See the Practice Considerations section for more information on populations at increased risk for acquiring STIs.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Skin Cancer Prevention: Behavioral Counseling: young adults, adolescents, children, and parents of young children	The USPSTF recommends counseling young adults, adolescents, children, and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons aged 6 months to 24 years with fair skin types to reduce their risk of skin cancer.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Statin Use for the Primary Prevention of Cardiovascular Disease in Adults: Preventive Medication: adults aged 40 to 75 years with no history of cvd, 1 or more cvd risk factors, and a calculated 10-year cvd event risk of 10% or greater	The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (ie, symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are aged 40 to 75 years; 2) they have 1 or more CVD risk factors (ie, dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults aged 40 to 75 years. See the "Clinical Considerations" section for more information on lipids screening and the assessment of cardiovascular risk.	Pharmacy Benefit	NA
Syphilis Infection in Nonpregnant Adults and Adolescents: Screening : asymptomatic, nonpregnant adults and adolescents who are at increased risk for syphilis infection	The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection.	86592-86593	Reason For Screen
Syphilis Infection in Pregnant Women: Screening: pregnant women	The USPSTF recommends early screening for syphilis infection in all pregnant women.	80055, 80081, 86592, 86593	Reason For Screen

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Unhealthy Alcohol Use in Adolescents and Adults: Screening and Behavioral Counseling Interventions: adults 18 years or older, including pregnant women	The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.	99408, 99409, G0396, G0397, G2011	Reason For Visit
Unhealthy Drug Use: Screening: adults age 18 years or older	The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.)	99408, 99409, G0396, G0397, G2011	Reason For Visit
Vision in Children Ages 6 Months to 5 Years: Screening: children aged 3 to 5 years	The USPSTF recommends vision screening at least once in all children aged 3 to 5 years to detect amblyopia or its risk factors.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral Interventions: adults	The USPSTF recommends that clinicians offer or refer adults with a body mass index (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive, multicomponent behavioral interventions.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit

Advisory Committee on Immunization Practices (ACIP) recommended vaccines

Follow current Medicaid policy for coverage of ACIP recommended vaccines including VFC.

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine

Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
Screening for gestational diabetes (Recommendation 5.1)	Screening for gestational diabetes in pregnant women between 24 and 28 weeks of gestation and at the first prenatal	82950-82952	See USPSTF A and B RECOMMENDATIONS Above

**Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines**

	visit for pregnant women identified to be at high risk for diabetes.		
--	--	--	--

**Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine
Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
Human papillomavirus testing (Recommendation 5.2)	The addition of high-risk human papillomavirus DNA testing in addition to cytology testing in women with normal cytology results. Screening should begin at 30 years of age and should occur no more frequently than every 3 years.	87623-87625, G0476	See USPSTF A and B RECOMMENDATIONS Above
Counseling for sexually transmitted infections (Recommendation 5.3)	Annual counseling on sexually transmitted infections for sexually active women.	Covered during preventive medicine services or a focused E/M visit.	See USPSTF A and B RECOMMENDATIONS Above
Counseling and screening for human immune-deficiency virus (Recommendation 5.4)	Counseling and screening for human immunodeficiency virus infection on an annual basis for sexually active women.	Counseling covered during preventive medicine services or a focused E/M visit. Labs as indicated. 86689, 86701-86703, 87389-87391, 87534-87539, 87806, G0432-G0433, G0435	See USPSTF A and B RECOMMENDATIONS Above
Contraceptive methods and counseling (Recommendation 5.5)	The full range of Food and Drug Administration-approved contraceptive methods, sterilization procedures, and patient education and counseling for women with reproductive capacity.	Counseling covered during preventive medicine services or a focused E/M visit. Oral contraceptives are included as a pharmacy benefit. 11976, 11981, 11982, 11983, 57170, 58300, 58301, 58565, 58600, 58605, 58611, 58615, 58670, 58671, A4264, A4266, A4268, A4269, J1050, J7296, J7297, J7298, J7300, J7301, J7304, J7307, S4989	The Medicaid program covers medically necessary contraceptive and counseling services for eligible beneficiaries.
Breastfeeding support, supplies and counseling (Recommendation 5.6)	Comprehensive lactation support and counseling and costs of renting breastfeeding equipment. A trained provider should provide counseling services to all pregnant women and to those in the postpartum period to ensure the successful initiation and duration of	A4281-A4286, E0602-E0604 S9443	See USPSTF A and B RECOMMENDATIONS Above

Michigan Department of Health and Human Services Preventive Services Coverage Guidelines

	breastfeeding. (The ACA ensures that breastfeeding counseling is covered; however, the committee recognizes that interpretation of this varies.)		
--	--	--	--

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
Screening and counseling for interpersonal and domestic violence (Recommendation 5.7)	Screening and counseling for interpersonal and domestic violence. Screening and counseling involve elicitation of information from women and adolescents about current and past violence and abuse in a culturally sensitive and supportive manner to address current health concerns about safety and other current or future health problems.	Covered during preventive medicine services or a focused E/M visit.	See USPSTF A and B RECOMMENDATIONS Above
Well-woman visit (Recommendation 5.8)	At least one well-women preventive care visit annually for adult women to obtain the recommended preventive services, including preconception and prenatal care. The committee also recognizes that several visits may be needed to obtain all necessary recommended preventive services, depending on a woman's health status, health needs, and other risk factors.	Covered during preventive medicine services.	Covered during preventive medicine services.

Early and Periodic Screening, Diagnosis and Treatment (EPSDT) services (19 and 20 year olds only)

Follow current Medicaid policy for coverage of EPSDT services.

Revision History

04/2014	<ul style="list-style-type: none"> - Added HCPCS codes: A4281-A4286 and E0602-E0604, to IOM's <i>Breastfeeding support, supplies and counseling preventive service.</i> - <i>Gestational diabetes mellitus screening</i> added to USPSTF January 2014
07/2014	<ul style="list-style-type: none"> - <i>Lung cancer screening</i> added to USPSTF December 2013 - <i>Hepatitis B screening: nonpregnant adolescents and adults</i> added to USPSTF May 2014 - <i>BRCA screening, counseling about</i> had an update to the USPSTF preventive service title. - <i>Dental caries prevention: preschool children</i> had an update to the USPSTF preventive service title. - Align <i>HIV screening</i> with current 2014 USPSTF version.
10/2014	<ul style="list-style-type: none"> - <i>Healthy diet counseling</i> had an update to the USPSTF preventive service title and description. - <i>Preeclampsia prevention: aspirin</i> added to USPSTF September 2014 - <i>Chlamydia screening: women</i> had an update to the USPSTF preventive service title and description. - <i>Gonorrhea screening: women</i> had an update to the USPSTF description. - <i>Sexually transmitted infections counseling</i> had an update to the USPSTF description.
02/2015	

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

	<ul style="list-style-type: none"> - Added 2015 HCPCS codes: 45346, 45388, 77063, 77085-77086, 87623-87625, 87806, G0464, G0472, G6019, G6022, G6024 - Deleted discontinued HCPCS codes: 45339, 45383, 87620-87622
10/2015	<ul style="list-style-type: none"> - High blood pressure in adults: screening added to USPSTF October 2015 - Diabetes Screening had an update to the USPSTF description.
01/2016	<ul style="list-style-type: none"> - Added HCPCS code: A4264 - Added 2016 HCPCS codes: 80081, 81528, G0297, J7297, J7298 - Deleted HCPCS code: G0464 as it is a Grade I. - Deleted discontinued HCPCS codes: G6019, G6022, G6024, J7302
04/2016	<ul style="list-style-type: none"> - Depression screening: adults had an update to the USPSTF preventive service description. - Depression screening: adolescents had an update to the USPSTF preventive service description. - Added HCPCS code: S9443 - Revised notes concerning Lactation support.
07/2016	<ul style="list-style-type: none"> - Diabetes screening: Added HCPCS codes: 82952 - Hepatitis B screening: nonpregnant adolescents and adults: Deleted HCPCS codes: 86707, 87341 - Hepatitis B screening: pregnant women: Deleted HCPCS codes: 80055, 80081, 86707, 87341 - Osteoporosis screening: women: Deleted HCPCS codes: 77085 and 77086. Added HCPCS codes: 76977, 77078, 77081 - Syphilis screening: nonpregnant/pregnant persons had an update to the USPSTF preventive service description. - Aspirin to prevent cardiovascular disease: men/women had an update to the USPSTF preventive service title and description. - Blood pressure screening in adults had an update to the USPSTF preventive service description. - Tobacco use counseling: nonpregnant/pregnant had an update to the USPSTF preventive service description. - Anemia screening: pregnant women removed from USPSTF A and B recommendation. - Iron supplementation in children: removed from USPSTF A and B recommendation. - Colorectal cancer screening had an update to the USPSTF preventive service description.
11/2016	<ul style="list-style-type: none"> - Tuberculosis screening: adults added to USPSTF September 2016 - Breastfeeding counseling had an update to the USPSTF preventive service title and description. - Statin preventive medication: adults ages 40–75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater added to USPSTF November 2016
01/2017	<ul style="list-style-type: none"> - Deleted discontinued HCPCS codes: G0436, G0437, 77052, 77057 - Added 2017 HCPCS codes: G0499, 77067 - Added HCPCS code: G0476
04/2017	<ul style="list-style-type: none"> - Cholesterol abnormalities screening: men 35 and older removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: men younger than 35 removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: women 45 and older removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: women younger than 45 removed from USPSTF A and B recommendation. - High blood pressure in adults: screening removed from USPSTF A and B recommendation.
07/2017	<ul style="list-style-type: none"> - Deleted HCPCS code: 99402 and note - Preeclampsia: screening added to USPSTF April 2017 - Obesity screening: children and adolescents had an update to the USPSTF preventive service title and description. - The format of this document was updated to include information reflective of the coverage of preventive services for both Healthy Michigan Plan beneficiaries and Medicaid Fee-for-Service beneficiaries.
09/2017	<ul style="list-style-type: none"> - Vision acuity screening: children had an update to the USPSTF preventive service title and description.
01/2018	<ul style="list-style-type: none"> - Alcohol misuse: screening and counseling: Added 99408 and 99409 - Deleted discontinued HCPCS codes: G0202, Q9984 - Added 2018 HCPCS codes: 00812, J7296
04/2018	<ul style="list-style-type: none"> - Skin cancer behavioral counseling: had an update to the USPSTF preventive service description. - Falls prevention: older adults: had an update to the USPSTF preventive service title and description. - Falls prevention in older adults: vitamin D removed from USPSTF A and B recommendation.
06/2018	

Michigan Department of Health and Human Services Preventive Services Coverage Guidelines

08/2018	- Osteoporosis screening: women: had an update to the USPSTF preventive service title and description.
09/2018	- Cervical cancer screening: had an update to the USPSTF preventive service description. - Syphilis screening: pregnant women: had an update to the USPSTF preventive service description. - Obesity screening and counseling: adults had an update to the USPSTF preventive service description.
10/2018	- Intimate partner violence screening: women of reproductive age had an update to the USPSTF preventive service title and description.
01/2019	- Alcohol misuse: screening and counseling had an update to the USPSTF preventive service title and description. - Gonorrhea prophylactic medication: newborns had an update to the USPSTF preventive service description.
02/2019	- Perinatal depression: counseling and interventions added to USPSTF February 2019.
06/2019	- HIV preexposure prophylaxis for the prevention of HIV infection added to USPSTF June 2019. - HIV screening: adolescents and adults ages 15 to 65 years had an update to the USPSTF preventive service title and description. - HIV screening: pregnant women had an update to the USPSTF preventive service description.
07/2019	- Hepatitis B screening: pregnant women had an update to the USPSTF preventive service description.
08/2019	- BRCA risk assessment and genetic counseling/testing had an update to the USPSTF preventive service description.
09/2019	- Breast cancer preventive medication had an update to the USPSTF preventive service description. - Bacteriuria screening: pregnant women had an update to the USPSTF preventive service description.
11/2019	- Added HCPCS codes: G0396, G0397, G2011
12/2019	- Abdominal aortic aneurysm screening: men had an update to the USPSTF preventive service description.
1/2020	- Added ICD-10 DX code: Z86.004
3/2020	- Hepatitis C virus infection screening: adults aged 18 to 79 years had an update to the USPSTF preventive service title and description.
4/2020	- Prevention and Cessation of Tobacco Use in Children and Adolescents: Primary Care Interventions: school-aged children and adolescents had an update to the USPSTF preventive service title and description.
6/2020	- USPSTF A and B Recommendation language updated to align with the USPSTF title and descriptions. - Unhealthy Drug Use: Screening: adults age 18 years or older added to USPSTF
8/2020	- Sexually Transmitted Infections: Behavioral Counseling: sexually active adolescents and adults at increased risk had an update to USPSTF preventive service title and description.
11/2020	- Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults With Cardiovascular Risk Factors: Behavioral Counseling Interventions: adults with cardiovascular disease risk factors had an update to USPSTF preventive service title and description.
12/2020	- Added HCPCS codes: 71271, 92650, 92651, 92652, 92653 - Deleted HCPCS code: 92585, 92586_G0297 - Screening for Hepatitis B Virus Infection in Adolescents and Adults: adolescents and adults at increased risk for infection had an update to the USPSTF preventive service title and description.
01/2021	- Interventions for Tobacco Smoking Cessation in Adults, Including Pregnant Persons: nonpregnant adults had an update to the USPSTF preventive service title and description.

Michigan Department of Health and Human Services Preventive Services Coverage Guidelines

	- Interventions for Tobacco Smoking Cessation in Adults, Including Pregnant Persons: pregnant persons had an update to the USPSTF preventive service title and description.
03/2021	- Lung Cancer: Screening had an update to the USPSTF preventive service title and description.
04/2021	- Screening for Hypertension in Adults: adults 18 years or older without known hypertension had an update to the USPSTF preventive service title and description.
05/2021	- Colorectal Cancer: Screening : had an update to the USPSTF preventive service title and description.